

ABSTRACT OF THE DISCLOSURE (New, Original)

Method of identification and quantitative analysis of carboxylic acid (s) in a sample by mass spectrometry using stable isotope labeled internal standard is provided. Said internal standard is prepared by reaction of an authentic sample of said carboxylic acid with a stable isotope labeled reagent, and is added to a sample containing said carboxylic acid. Said carboxylic acid in said sample is then quantitatively converted to a chemical compound of identical structure, except the stable isotope atoms, as that of said internal standard using a non-labeled reagent. Said sample is then extracted and the extract is analyzed by mass spectrometry. Identification and quantification of said carboxylic acid are made from a plot of ion ratio of said converted carboxylic acid to said internal standard versus carboxylic acid concentration.

REFERENCES (Original)

US patent documents

5,559,038	September 24, 1996	J. Fred Kolhouse
6,358,996	March 19, 2002	Michael S. Alexander

Other references

Petr Husek, "Chloroformates in gas chromatography as general purpose derivatizing agents", Journal of Chromatography B, 1998, page 57-91, vol. 717.

Jens Pietzsch et al, "Rapid determination of total homocysteine in human plasma by using N(O,S)-Ethoxycarbonyl ethyl ester derivatives and gas chromatography-mass spectrometry", Clinical Chemistry, 1997, page 2001-2004, vol. 43(10).

Petr Husek and Petr Simek, "Advances in amino acid analysis", LCGC Sep 2001, page 986-999, vol.19.

Ping Cao and Mehdi Moini, "Quantitative analysis of fluorinated ethylchloroformate derivatives of protein amino acids and hydrolysis products of small peptides using chemical ionization gas chromatography-mass spectrometry", Journal of Chromatography A, 1997, page 111-117, vol. 759.

William A. Joern, "Unexpected volatility of barbiturate derivatives : an extractive alkylation procedure for barbiturates and benzoylecgonine", Journal of Analytical Toxicology, Nov 1994 page 423, vol. 18.